*Grip*Protect® Precise BLACK 5 Nitrile Exam Gloves

Formulated to provide superior barrier protection



*Grip*Protect Precise BLACK 5 Nitrile Exam Gloves offer durable barrier protection in many industries ranging from healthcare to industrial and laboratory environments. Thicker than our standard 3.5 mil GripProtect Precise BLACK Nitrile gloves, the BLACK 5 gloves have a comfortable and durable feel with better protection.

Medical Exam Grade classification in accordance with ASTM testing criteria ensures the gloves perform consistently and to the highest accepted standards to protect the hands of the users and the people on the other side. Our gloves exhibit a similar feel, flexibility and tactile sensitivity as latex gloves, but contain no natural rubber latex so they are safe for people with latex allergies.

- Medical exam grade
- Chemo rated*
- Latex-free, Powder-Free
- Non-sterile
- Ambidextrous, black color
- Textured fingers



* Chemotherapy drug testing using the ASTM D 6978-05 permeation test. See Chemotherapy Drug Resistance data sheet for complete testing details.



For more information visit: www.bmcprotect.com 1-800-977-7888

*Grip*Protect® Precise BLACK 5 Nitrile Exam Gloves

- Medical exam grade
- Chemo rated
- Latex-free, Powder-Free
- Non-sterile
- Ambidextrous
- Textured fingers



Part Number	Size	# Gloves	Weight / Glove (g)	Width at Palm (cm)
GP2451	Small	100/Box, 1000/Case	3.97 ± 0.2	8.5 ± 1
GP2452	Medium	100/Box, 1000/Case	4.62 ± 0.2	9.5 ± 1
GP2453	Large	100/Box, 1000/Case	4.94 ± 0.2	11.0 ± 1
GP2454	Extra-Large	100/Box, 1000/Case	5.16 ± 0.2	11.5 ± 1

Specifications, Physical Properties, & Quality Standards					
Characteristic	Standard	Test Method			
Tensile Strength before aging Tensile Strength after aging	≥ 14 Mpa ≥ 14 MPa	ASTM D412/ASTM D573 ASTM D412/ASTM D573			
Elongation before aging Elongation after aging	≥ 500% ≥ 400%	ASTM D412/ASTM D573 ASTM D412/ASTM D573			
Thickness at middle finger Thickness at palm	0.11 mm (4.33 mil) 0.08 mm (3.15 mil)	ASTM D3767 ASTM D3767			
Barrier properties	AQL 1.5	ASTM 5151			
Length	> 240 mm	ASTM D3767			



*Grip*Protect® Precise BLACK Nitrile Exam Gloves



Chemotherapy Drug Resistance

Tested for use with Chemotherapy drugs using the ASTM D 6978-05 permeation test. No breakthrough after 240 minutes for the below drugs at the listed concentration.

Test Chemotherapy Drug	Drug Concentration	
5-Fluorouracil (Adrucil)	50.0 mg/ml (50,000 ppm)	
Etoposide (Toposar)	20.0 mg/ml (20,000 ppm)	
Cyclophosphamide (Cytoxan)	20.0 mg/ml (20,000 ppm)	
Paclitaxel (Taxol)	6.0 mg/ml (6,000 ppm)	
Doxorubicin Hydrochloride (Adriamycin)	2.0 mg/ml (2,000 ppm)	
Dacarbazine	10.0 mg/ml (10,000 ppm)	
Cisplatinum	1.0 mg/ml (1,000 ppm)	
Ifosfamide	50.0 mg/ml (50,000 ppm)	
Mitoxantrone	2.0 mg/ml (2,000 ppm)	
Vincristine Sulfate	1.0 mg/ml (1,000 ppm)	
Reduced permeation time compounds.		
Carmustine (BCNU) – 15 minutes	3.3 mg/ml (3,300 ppm)	
Thiotepa – 2 minutes	10.0 mg/ml (10,000 ppm)	

